


# Fundamental theorem of calculus

Provided that the integrand is continuous on  $[a, b]$ , the **integral undoes the derivative**

$$\int_a^b \frac{df}{dx} dx = f(b) - f(a)$$

$f$  is an “antiderivative” of  $\frac{df}{dx}$ .

“The integral over an interval is equivalent to a function evaluated at the bounds.”




Provided that the integrand is continuous on  $[a, b]$  and that  $x \in [a, b]$ , the **derivative undoes the integral**

$$\frac{d}{dx} \underbrace{\int_a^x f(t) dt}_{A(x)} = f(x)$$

“The rate at which area is accumulated is the function.”

$$\begin{aligned} \frac{dA}{dx} &= \lim_{\Delta x \rightarrow 0} \frac{A(x + \Delta x) - A(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{f(x)\Delta x + \mathcal{O}(\Delta x^2)}{\Delta x} \\ &= f(x) \end{aligned}$$



## Antiderivative

Because the integral is the inverse of the derivative, we can reverse differentiation formulas to obtain integration formulas.

*Example:* Power rule

$$\frac{d}{dx} [x^n + \text{any constant}] = nx^{n-1}$$

implies

$$\int_a^b x^n dx = \left[ \frac{x^{n+1}}{n+1} + C \right]_a^b, \quad n \neq -1$$

When the  $\int$  and  $dx$  appear without limits, they can together be read as “antiderivative” of whatever they surround.

We abuse the  $\int$  and  $dx$  symbols in the same way that we abuse symbols that denote derivatives. We apply the  $\int$  and  $dx$  both (1) to names of functions and (2) to association rule formulas in terms of  $x$ .

Even though antiderivative identities can be derived from differentiation formulas, memorize a basic set of antidifferentiation identities (see separate table).

## u-substitution

$$\int_{x=1}^{x=2} (x+1)^2 dx$$

$$u = x + 1 \quad x = 1: u = (1) + 1 = 2$$

$$du = (1 + 0)dx \quad x = 2: u = (2) + 1 = 3$$

$$\int_{u=2}^{u=3} u^2 du$$